

# Towards a conceptual framework:

Trauma, Family Violence and Health

---



## Suggested Citation:

Baker, L., Straatman, A.L., Etherington, N., O'Neil, B., Heron, C., Sapardanis, K. (2016). Towards a conceptual framework: Trauma, Family Violence and Health. London, ON: Knowledge Hub, Learning Network, Centre for Research & Education on Violence against Women & Children.

ISBN 978-1-988412-09-2

Copyright: Centre for Research & Education on Violence against Women & Children




**Western**

Centre for Research & Education on  
Violence Against Women & Children

# Contents

<b>Trauma, Family Violence, and Health</b> .....	4
Introduction .....	4
Defining Trauma in the Context of IPV and Child Maltreatment .....	4
Considering the Health Impacts of Trauma.....	8
Toward a Framework for Understanding Violence, Trauma and Health.....	11
Socio-ecological Model .....	11
The Life Course Perspective.....	11
Intersectionality: Bringing context to the center .....	12
A Trauma-Informed Health Framework.....	13
<b>Conclusion</b> .....	15
<b>References</b> .....	16



# Trauma, Family Violence, and Health

## Introduction

Family violence is a public health issue with pervasive consequences, including experiences of trauma. Trauma has been linked to a variety of short- and long-term physical and mental health outcomes, including sleep problems, weakened immune system functioning, diabetes, hypertension, cardiovascular disease, post-traumatic stress disorder (PTSD), anxiety, and depression (Lind et al., 2016; Petrov et al., 2016; Lagdon, Armour, & Stringer, 2014; Mason et al., 2013; Out et al., 2012; Irish et al., 2010; Dutton et al., 2006; Seng et al., 2005; Woods et al., 2005; Reiche, Nunes & Morimoto, 2004; Shaw & Krause, 2002; Clum et al., 2001). From a public health perspective, increased attention has been directed to trauma resulting from family violence and its effects on well-being. Continued efforts to bring the field of public health and trauma-informed responses to survivors of family violence is integral to advancing prevention and intervention across the life course, as well as improving services and health for survivors of violence.

Broadly, family violence encompasses all forms of abuse or neglect experienced by a child or adult from a family member or person with “whom they have an intimate relationship” (Public Health Agency of Canada, 2014). Examples of family violence

identified by the Public Health Agency of Canada (2014) include: elder abuse/neglect, early and forced marriage, violence directed toward women and girls in the name of “honour”, female genital mutilation, intimate partner violence, and child maltreatment (including abuse, neglect and exposure to IPV). This paper focuses on intimate partner violence (IPV) and child maltreatment. Both IPV and child maltreatment are highly prevalent forms of family violence worldwide and represent serious public health concerns, with a range of negative impacts on well-being across the life course.

We begin by reviewing the health outcomes of trauma within the context of IPV and child maltreatment. Next, three theoretical models and their relevance to trauma-informed health promotion for child and adult survivors are presented. Finally, incorporating the model of SAMHSA (SAMHSA, 2014a), we describe a trauma-informed health framework that will provide the foundation for the development of outcome indicators for trauma-informed health promotion.

## Defining Trauma in the Context of IPV and Child Maltreatment

Research indicates that approximately 76% of adults in Canada report having been exposed to at least some form of trauma in their lifetime, with about 9% meeting the criteria for post-traumatic stress disorder (PTSD) (Van Ameringen, 2008). Trauma can result from a single event, series of events, or set of circumstances which threaten(s) a person’s physical or psychological well-

being and overwhelm(s) their capacity to cope (BC Provincial Mental Health and Substance Use Planning Council, 2013). Experiences of trauma vary in magnitude, frequency, duration, complexity, and source. Along these dimensions, there are five types of trauma, summarized in the chart below.

**Table 1. Types of Trauma**

TYPE	DESCRIPTION	EXAMPLES
<b>Single incident trauma</b>	Trauma related to an unexpected and overwhelming event	Car accident, natural disasters, sudden loss, community violence (e.g. school shooting), single episode of abuse or assault, witnessing violence
<b>Complex or repetitive trauma</b>	Trauma related to ongoing traumatic experiences	War, immigrant/refugee experiences (e.g. forced displacement), ongoing abuse, intimate partner violence
<b>Developmental trauma</b>	Trauma related to early ongoing or repetitive trauma, often within a child's caregiving system and interfering with healthy attachment and development	Neglect, abandonment, physical abuse or assault, sexual abuse or assault, emotional abuse, witnessing violence or death, coercion or betrayal
<b>Intergenerational trauma</b>	Psychological or emotional effects that can be experienced from people who live with trauma survivors	Children living with a parent or caregiver who experienced abuse, children of survivors of residential schools
<b>Historical trauma</b>	Cumulative emotional and psychological wounding over the lifespan and across generations emanating from massive group trauma, inflicted by a subjugating, dominant population; intergenerational trauma is an aspect of historical trauma	Genocide, colonialism, slavery, war

Source: BC Provincial Mental Health and Substance Use Planning Council, 2013.

While there are many experiences which can result in trauma, we focus on IPV and child maltreatment – which are examples of, or strongly related to, each of the aforementioned types. It is important to note that IPV and child maltreatment do not always lead to trauma. Although many violent experiences are inherently considered to be traumatic, the event itself is not the determining factor of whether someone experiences trauma or not; rather, “it is the individual’s experience of the event and the meaning they assign to it” (Klinik Community Health Centre, 2013, p.9). That is, two individuals may be exposed to the same event, yet differ on what is interpreted as traumatic (SAMHSA, 2014). Protective factors (e.g. self-esteem, social support, self-regulation) following a potentially traumatic event can negate the preceding trauma; however, not all individuals have the capacity for or are able to access these protective factors. As a result, some individuals may have a greater risk of experiencing

consequences to health and well-being following trauma (Klinik Community Health Centre, 2013; Pearlin, 2010). This paper centers on instances where IPV and child maltreatment results in trauma, resulting in both direct and indirect effects on health. We also recognize that survivors of IPV and child maltreatment may have previously experienced or may continue to experience other forms of trauma beyond family violence (e.g. a survivor of intimate partner violence may also be affected by intergenerational trauma related to residential schools).

Intimate partner violence refers to a range of abusive behaviours perpetrated by a current or former partner, including but not limited to: physical, sexual, and psychological or emotional harm (Sinha, 2013). These dimensions of IPV are defined in Table 2.

“Individual trauma results from an event, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful or life threatening and that has lasting adverse effects on the individual’s functioning and mental, physical, social, emotional, or spiritual well-being.”  
SAMHSA, 2014b, p.7.

## Table 2. Forms of IPV and definitions

<b>Physical Violence</b>	Physical injury or death resulting from the intentional use of physical force (e.g. pushing, shoving, throwing, biting, hair-pulling, choking, slapping, punching, burning, use of a weapon, use of one's body, size or strength against another person)
<b>Sexual Violence</b>	Sexual acts committed against an intimate partner without freely given consent (e.g. physical force or alcohol/drug induced sexual acts, "non-physical" pressure to engage in sexual acts—through fear, verbal pressure, intimidation, forcing a partner to engage in unsafe or humiliating sexual acts, and non-contact acts of a sexual nature—sexual comments, voyeurism, exhibitionism)
<b>Psychological or Emotional Violence</b>	The use of words or action is to control or frighten an intimate partner, or destroy their self-respect (e.g. insults, humiliation in front of others, use of intimidation, coercive control, hurting pets, destroying belongings, threats of physical or sexual violence, control of reproductive or sexual health—hiding birth control pills, exploitation)
Source: Adapted from Sinha, 2013 & PHAC, 2014.	

IPV has particular impacts on the formation of trauma. Specifically, IPV is typically chronic in nature. It is often ongoing yet unpredictable, leading to more powerful, negative consequences (Katerndahl et al., 2010; Engnes, Liden & Lundgren, 2012; Herman, 1992a,b). Such violence is also perpetrated by a loved one, leading to a violation of trust and disruption of attachment, which can be highly traumatic itself (Scott & Babcock, 2010).

IPV can occur in all types of relationships, regardless of the gender of the individuals involved in the relationship, which may or may not include sexual intimacy. Police-reported Canadian data indicate that there were 341 victims per 100,000 population of intimate partner violence in 2011, with 80% of victims being women (Sinha, 2013). Rates are particularly high for young women (age 15 to 24 and 25 to 34) and marginalized women (e.g. Indigenous women). At a global level, approximately 30% of women have been subjected to physical and/or sexual violence by an intimate partner (WHO, 2013). Though IPV is most commonly experienced by women and perpetrated by men, men can also be survivors of violence. Police-reported data indicates that 147 men per 100,000 have experienced intimate partner violence in

Canada, compared to the rate for women at 574 per 100,000 (Sinha, 2012). Rates for men may be underestimated, however, given US findings that men are less likely than women to report experiences of violence (Black et al., 2011).

Child maltreatment includes all forms of physical, sexual, and psychological abuse directed toward a child as well as neglect and exposure to IPV (Public Health Agency of Canada, 2010). Specific forms of child maltreatment are further defined with examples in Table 3.

### Table 3. Forms of child maltreatment and definitions

<b>Physical abuse</b>	Any act of physical aggression directed toward a child (e.g. shaking, pushing, hitting with object, biting, choking).
<b>Sexual abuse</b>	Sexual molestation or exploitation of a child by an adult or older child within or outside the family (e.g. penetration, fondling, pornography).
<b>Emotional/psychological abuse</b>	Terrorizing or threat of violence (e.g. threats against child's cherished objects), verbal abuse or belittling (e.g. name-calling), isolation or confinement (e.g. purposely cutting child off from other children), inadequate nurturing or affection (e.g. lack of parental interaction), exploiting or corrupting behaviour (e.g. encouraging involvement in criminal behaviour).
<b>Neglect</b>	Failure to provide for child's basic needs, adequate protection, and adequate supervision (e.g. inadequate nutrition, failure to provide medical treatment).
<b>Exposure to intimate partner violence</b>	Child is present during physical or verbal violence between intimate partners and can see and/or hear the violence (direct); child not present during violence but suffers consequences, hears about it, or experiences changes in his/her life as a result (indirect); child is exposed to emotional violence between intimate partners.
Source: Public Health Agency of Canada, 2010.	

Cases of child maltreatment are difficult to identify and rates of child maltreatment can vary depending on the methods used to gather data (Hambrick et al., 2014). The Canadian Incidence Study of Child Maltreatment (CIS) is one of the few nationwide studies to investigate the prevalence of child maltreatment cases (Tonmyr, Ouimet, & Ugnat, 2012). This study utilized direct reports from child welfare workers to uncover maltreatment rates for the four categories of maltreatment. Of the 6,163 substantiated investigations of maltreatment among children of all ages, the primary category of maltreatment was physical abuse in 20% of cases, sexual abuse in 3% of cases, emotional abuse in 9% of cases, neglect in 34% of cases, and exposure to IPV in 34% of cases (PHAC, 2010). Further analysis of the data from this study focusing on First Nations children found that of the substantiated cases of maltreatment, the primary category of maltreatment was neglect in 46% of cases, exposure to intimate partner violence in 36% of cases, physical abuse in 23% of cases, emotional maltreatment in 9% of cases, and sexual abuse in 3% of cases (Sinha, Trocme, Fallon, MacLaurin, Fast, Prokop et al, 2011).

The consequences of child maltreatment have direct impacts on healthy development as well as achievement, which can increase the likelihood of disease in adulthood (WHO, 2016). In fact, the World Health Organization identifies child maltreatment as a contributing factor to slowed economic and social development at the nation-level. Furthermore, the poor health outcomes resulting from child maltreatment form a significant portion of the global burden of disease (WHO, 2016). It is also likely to occur in the context of IPV, with prevalence estimates for co-occurrence ranging from 30 to 60% (Edleson, 1999; Jouriles et al., 2008; Hamby et al., 2010) and can be a risk factor for the future perpetration of IPV (see Etherington & Baker, 2016a, for a review). Child maltreatment can be particularly traumatic because it is perpetrated by a caregiver - a person whom the child depends on for their basic needs and protection in addition to love, warmth, and support. This type of family violence can also be chronic in nature, often involving multiple forms of abuse, which can further exacerbate experiences of trauma.

# Considering the Health Impacts of Trauma

## Social Determinants of Health

The following factors have been shown to strongly effect the health of Canadians:

- Aboriginal Status
- Disability
- Early life
- Education
- Employment and working conditions
- Food insecurity
- Health services
- Gender
- Housing
- Income and income distribution
- Race
- Social exclusion
- Social safety net
- Unemployment and job security

Raphael (2009)

Substantial research has documented the impacts of trauma and family violence on health and well-being (e.g. Lum et al., 2016; Lopez-Martinez et al., 2016; Ely et al., 2004; Lagdon et al., 2014; Macy et al., 2009; Wong & Mellor, 2014; WHO, 2013; Blasco-Ros, Herbert & Martinez, 2014; Herrenkohl et al., 2013; Dillon et al., 2013). This section summarizes the physical, psychological, behavioural and interpersonal health consequences found to be specifically related to trauma that occurs in the context of IPV and child maltreatment (see Table 4). In general, the health consequences of trauma are more severe with multiple or chronic exposures and can be long-lasting, particularly with regard to mental health (for a review, see D'Andrea

et al., 2011). Trauma emerging from IPV or child maltreatment can have a variety of direct and indirect health consequences, as well as the potential to lead to further co-occurring disorders and/or concerns, such as substance use.

The health impacts of trauma should be considered in the context of pre-existing health inequalities or differences in health among Canadians based on the structural circumstances, or social and economic conditions, in which individuals live. Research demonstrates that factors such as gender, income, education, ethnicity/culture, Aboriginal status, and immigrant status are key factors in shaping health outcomes (Health Disparities Task Group, 2005; Canadian Population Health Initiative, 2004; Government of Canada, 2008; Canadian Population Health Initiative,

2008). These and other factors, commonly referred to as the social determinants of health (see PHAC, 2016) are associated with inequalities in such outcomes as mortality, early childhood development, health behaviours (e.g. smoking, exercising), and health care utilization (e.g. visits to the doctor), mental illness, morbidity, and disability (Pan-Canadian Public Health Network, 2010). Thus, some individuals may have an increased vulnerability to health problems, which can subsequently affect or be affected by their trauma experience.

Due to the prolonged, re-occurring trauma associated with IPV, there is an increasing likelihood of developing severe health consequences (e.g., Lum et al., 2016; Beydoun, Beydoun, Kaufman, Lo, & Zonderman, 2012; Ely, Dulmus, & Wodarski, 2004; Haskell, 2001; Jones, Hughes, & Unterstaller, 2001; Lagdon et al., 2014; Macy, Ferron, & Crosby, 2009; Wathen, 2012; Wong & Mellor, 2014; WHO, 2013). The most prevalent health impacts of IPV are PTSD and depression. Rates of PTSD in women who have experienced IPV have been shown to range in magnitude, from 14% to 92% of women exhibiting at least some symptoms (Dillon et al., 2012). PTSD has also been found to be a long-lasting effect of IPV, with some women experiencing high rates of PTSD symptoms as many as nine years after an abusive relationship has ended (Wong & Mellor, 2014). With regard to depression, women who have experienced IPV victimization appear to experience rates two to five times higher than women without such an experience (Beydoun et al., 2012; Wathen, 2012). While the type of IPV victimization has not been found to play a large role in the development of depression, it has been suggested that psychological abuse alone may lead to even higher rates of depressive symptoms (Lagdon et al., 2014). In addition, repeated victimization greatly increases the risk of developing major depression (Lagdon et al., 2014).

The physical health impacts of IPV are far-ranging, from short-term injuries to death. Such consequences of violence have been divided into three distinct categories: immediate and direct impacts (i.e., death and injury); long-term and direct impacts (i.e., disability and chronic illness); and indirect impacts (i.e., health behaviours,

such as smoking, diet, and physical activity) (Wong & Mellor, 2014). Although some consequences are more serious than others, all physical health impacts resulting from IPV can be life altering. Table 4 contains a summary of the effects of IPV on the health and well-being of survivors.

**Table 4. Effects of IPV on health and well-being.**

PHYSICAL	PSYCHOLOGICAL/ EMOTIONAL	BEHAVIOURAL	INTERPERSONAL
Physical injuries (e.g. cuts, bruises, sprains, broken or fractured bones)	Post-traumatic stress disorder	Eating disorders	Frequent relationship conflict
Brain injury	Anger management	Substance abuse	Experiences of re-victimization
Cardiovascular disease	Anxiety	High medication use	Perpetration of violence
Hypertension	Depressive symptoms	Self-harm	Difficulty establishing and maintaining relationships
Arthritis	Major depressive disorder	High-risk sexual behaviours	
Irritable bowel syndrome	Suicidality		
Chronic pain	Obsessive-compulsive disorder		
Reproductive and gynecological health problems	Poor self-rated mental wellness		
Somatoform symptoms	Poor emotional regulation		
Poor self-reported physical health and quality of life			
Sleep problems			
Digestive problems			
Disability			
Death			

Adapted from: Lum et al., 2016; Wong & Mellor, 2014; WHO, 2013.

Research has also consistently linked child maltreatment to numerous health outcomes that can span across the life course (e.g. Briere & Jordan, 2009; Fry et al., 2012; Hébert et al., 2016). One examination of substantiated cases of maltreatment found that 28% of children experienced health problems other than immediate physical injury, such as asthma or non-organic failure to thrive (Health Canada, 2005). Emotional harm appears to be more common than physical harm, with 34% of children experiencing emotional harm, and 21% requiring treatment. This outcome was particularly prevalent for children who experienced sexual abuse (Health Canada, 2005).

Recent Canadian research indicates that these negative health outcomes often continue into adulthood for children who have experienced maltreatment. Specifically, findings indicate increased odds of physical conditions (e.g. back problems, cancer, chronic fatigue, stroke), poor self-perceived health, and mental health issues (e.g. anxiety, depression, and post-traumatic stress disorder) (Afifi et al., 2016; Afifi et al., 2014). In many instances, the odds of experiencing a physical condition in adulthood were approximately 1.5 to 3 times greater compared to individuals who had not experienced child abuse (Afifi et al., 2016). Notably, odds ranged from 1.4 to 7.9 for mental health issues (Afifi et al., 2014).

In other words, victims of childhood abuse are approximately 1.5 to 8 times more likely to experience psychological or emotional consequences in adulthood, depending on the outcome examined. Suicidality is of particular concern given findings

that experiences of sexual abuse result in an increased likelihood of attempting suicide 8 times that of individuals without such an experience. The effects of child maltreatment on the health and well-being of children are summarized in Table 5.

**Table 5. Effects of child maltreatment on health and well-being**

PHYSICAL	PSYCHOLOGICAL/ EMOTIONAL	BEHAVIOURAL	INTERPERSONAL
Pain Increased number of surgeries Increased rates of hospitalization Somatic symptoms Obesity Chronic illness/disease in adulthood/after life	Anxiety Major depressive disorder Oppositional-defiant disorder Conduct disorder Post-traumatic stress disorder Psychological distress Low self-esteem Suicidality Psychiatric symptoms Antisocial behaviour & personality disorders Emotion regulation difficulties Cognitive disturbances	High-risk sexual behaviours Early initiation of smoking Substance use and abuse (drugs and alcohol) Eating disorders Self-harm Educational impacts (poor academic achievement)	Vulnerable to later re-victimization Difficulty forming relationships Dating violence/IPV perpetration Bullying
Briere & Jordan, 2009; Fry et al., 2012; Rodgers et al., 2004; Zimmerman & Mercy, 2010; Rogosch et al., 2011; Herrenkohl et al., 2008; Romano, Babchishin, Marquis, & Fréchette, 2015.			

# Toward a Framework for Understanding Violence, Trauma and Health

The application of the social-ecological model to trauma and its effects as well as to public health prevention can be complemented by other existing health frameworks, such as the

## Socio-ecological Model

The socio-ecological model has been fundamental to public health's approach to violence prevention (for examples see, CDC, 2015; Miskurka, Steensma, & Phillips, 2016). This model views violence as a complex behaviour that is best explained by the interactions of multiple factors related to individuals and the contexts within which they live, learn, play, and work. Specifically, four contexts are identified as levels of influence contributing to violence and as sites to stop violence, to respond to those affected or at risk of being affected by violence, and to prevent violence from beginning: individual, relationship, community, and societal. Table 6 summarizes the factors associated with each level of the model.

Risk and protective factors at each of the levels are targeted through prevention and intervention initiatives. There are thus

## The Life Course Perspective

A life course approach to health emphasizes the temporal and social dimensions of health. Within this perspective, health is viewed as a dynamic process extending from the prenatal period through to old age. This process is influenced by social, economic, and cultural contexts and occurs not only across individuals' lives, but also, across generations. Further, health and the factors that shape it are not static. In other words, health and its determinants can change over time. For example, an individual may be unemployed for several years and then employed again, and in turn, experience benefits to health that they formerly did not possess.

life course perspective (see Elder, Johnson & Crosnoe, 2003) and intersectionality theory (see Hankivsky, 2011).

numerous opportunities for implementing a multifaceted, multi-level prevention strategy. For example, health promotion for trauma survivors at the individual level may include educating high school students about health risk behaviours associated with trauma/family violence (e.g. smoking, drinking, drugs) and include programming for alternative health-promoting coping strategies (e.g. yoga, sports). At the interpersonal level, students might participate in healthy relationships programming that teaches about negotiating safe-sex practices with a partner. Community-level prevention might include a youth health drop-in centre offering confidential assessment, STI-testing, and counselling services. Finally, societal initiatives might include policies to provide funding for low-income adolescents to participate in team sports.

In its recognition of dynamic processes, the life course perspective emphasizes the link between early life events and later life outcomes, and the cumulative impact of these events. This is an important tenet to consider given the often cumulative nature of violence, trauma, and its related stressors. For example, a survivor may have had experiences of physical abuse during childhood by a parent, sexual abuse during adolescence by a peer, and emotional abuse during a marital relationship. Apart from re-victimization, violence can have a cumulative effect within one relationship as it is often a process rather than an event (Williams, 2003). IPV, for example, is often chronic and enduring, with each episode of violence building on previous episodes.

**Table 6. Socio-ecological Model Levels and Factors**

LEVELS:	FACTORS:
<b>Individual</b>	Identifies biological and personal history factors; such as age, education, income, substance use, or history of abuse, that increase the likelihood of becoming a victim or perpetrator of violence
<b>Relationship</b>	Examines close relationships that may increase the risk of experiencing violence as a victim or perpetrator. A person’s closest social circle-peers, partners and family members-influences their behavior and contributes to their range of experience.
<b>Community</b>	Explores the settings, such as schools, workplaces, and neighborhoods, in which social relationships occur and seeks to identify the characteristics of these settings that are associated with becoming victims or perpetrators of violence.
<b>Societal</b>	Looks at the broad societal factors, such as health, economic, educational and social policies, that help create a climate in which violence is encouraged or inhibited and help to maintain economic or social inequalities between groups in society.
Source: CDC., (2015) The Social-Ecological Model	

## Intersectionality: Bringing context to the center

Intimate partner and sexual violence scholarship and activism are grounded in an intersectional understanding of survivors’ vulnerabilities to and experiences with violence, in addition to the systems and services with which they become involved (e.g. Sokoloff, 2005; Crenshaw, 1991). Intersectionality draws attention to the ways in which lived experiences and life changes are shaped by interconnected dimensions of stratification (e.g. ability, indigeneity, gender, age) as well as the broader social context (e.g. social disadvantage, historical and current oppressions). Accordingly, not all survivors of family violence experience IPV or child maltreatment in the same way, and their experiences of trauma and health occur in – and are impacted by – different contexts.

When research examines violence, trauma, and health among various groups, it often fragments individuals into single or dyadic categories. In reality, individuals may identify as belonging to several different groups and can be impacted by the multiple oppressions facing each group. In addition, no single axes of inequality is more important than any other as these axes are

interconnected and work simultaneously to shape individuals’ experiences.

It is important to note that while occupying multiple disadvantage statuses can have implications for individual outcomes, these implications are not additive in nature (Dill & Zambrana, 2009). That is, “the impact of intersecting identities is qualitatively different from the impact of any single identity or the addition of them” (Etherington & Baker, 2016b, p. 3, emphasis added). Survivors of violence navigate multiple social statuses and experiences of oppression simultaneously, and have unique experiences of trauma and health as a result. It is not a question of whether one person experiences “greater” or “worse” effects than another; rather, it is a matter of understanding the context in which these experiences occur. Further, when interpreting available research on violence, trauma, and health among diverse groups, it is important to avoid conflating experiences of violence with stereotypical accounts of particular groups as this may overlook the complexities of lived experience and reinforce oppressive discourses (e.g. racism, ableism, homophobia, transphobia).

From a trauma-informed lens, it is critical that services avoid re-traumatizing individuals and the risk of re-traumatization is greater for those experiencing multiple forms of oppression (Miller et al., 2016). There has also been increasing recognition of the need for trauma researchers and practitioners to consider

the diverse experiences of survivors (see Bryant-Davies, 2010). Intersectionality is therefore a key element of trauma-informed approaches to improve the health and well-being of survivors of violence given the importance of context for these outcomes.

## A Trauma-Informed Health Framework


A social-ecological model (SEM) informed by the life course perspective and intersectionality can be applied to both trauma and health, as it involves many of the same elements in each instance while recognizing the reciprocity of each level (i.e. individual, interpersonal, community, societal) in shaping relevant outcomes. Accordingly, it makes sense to integrate approaches to trauma and health through the social-ecological model, which has also been fundamental to intervention strategies (see, for example, CDC, 2015; Magruder et al., 2016). Such a model has been developed by SAMHSA (2014a) and is presented in Figure 1.

The SEM model provides a strong conceptual base for trauma-informed health promotion. Specifically, trauma is viewed through a broad lens that incorporates individual,

interpersonal, community, and societal level factors, all of which are present before, during and after the trauma. It recognizes that environmental factors, including our access to resources, greatly influence all aspects of our well-being (see Social Determinants of Health, on p.8). The framework supports primary, secondary and tertiary prevention initiatives that work through inter-related factors at each of the social-ecological levels while acknowledging the link between early and later experiences and how this may be shaped by intersecting social inequalities (see Table 7). When applied to health promotion, this model brings trauma to the forefront of developmentally sensitive intervention efforts and health outcomes.

Figure 1. A Social-Ecological Model for Understanding Trauma and Its Effects

## Trauma-Informed Health Framework for Survivors of Family Violence



Source: SAMSHA, 2014, Page 36

**Table 7. Examples of factors operating at each level within the social-ecological model of trauma and its effects.**

INDIVIDUAL FACTORS	INTERPERSONAL FACTORS	COMMUNITY AND ORGANIZATIONAL FACTORS	SOCIETAL FACTORS	CULTURAL AND DEVELOPMENTAL FACTORS	PERIOD OF TIME IN HISTORY
Age Biophysical state Mental health status Personality traits (e.g. temperament) Education Gender Coping styles Socioeconomic status	Family Peer and significant other interaction patterns Parent/family mental health Parents' history of trauma Social network	Neighbourhood quality School system and/or work environment Social services, family violence services, and health services quality and accessibility Faith-based settings Transportation availability Community socioeconomic status Community employment rates	Laws Provincial and Federal economic and social policies Media Societal norms Judicial system	Collective or individualistic cultural norms Ethnicity Cultural subsystem norms Cognitive and maturational development	Societal attitudes related to family violence Changes in diagnostic understanding between editions of the Diagnostic and Statistical Manual of Mental Disorders
*List not exhaustive. Adapted from SAMHSA, 2014a, p. 16.					

## Conclusion

The burden of victimization is evident in the emerging body of evidence linking experiences of family violence as children and/or as adults to trauma, and in turn, to poor health outcomes. While not all experiences of violence lead to trauma and not all trauma results in poor health, a substantial number of survivors experiencing trauma face resulting physical, psychological/emotional, behavioural, and social hardships (Lum et al., 2016; WHO, 2013; Wong & Mellor, 2014). The fields of public health and trauma-informed approaches are foundational to supporting survivors of IPV and child maltreatment.

Socio-ecological, life course, and intersectional approaches contribute explanatory value to a trauma-informed health framework. The integration of these frameworks recognizes the social-ecological influences on trauma, and in turn, on health, while at the same time recognizing temporal patterns and intersectional experiences. Key to such a framework are the following premises: there is a link between and cumulative impacts of events

in early life and later life outcomes; risks and protectors linked to violence are associated with individual factors (e.g. biological factors) and contextual factors (e.g. relationship, community, societal factors); experience and life changes—including experiences of violence, trauma and help seeking—are shaped by interconnected dimensions of stratification (e.g. ability, indigeneity, gender, age) as well as the broader social context (e.g. social disadvantage, historical and current oppressions); and, survivors of family violence experience IPV or child maltreatment in different ways.

The elements of the framework described above are represented in the social-ecological model for understanding trauma and its effects developed by SAMHSA (2014a). This heuristic framework provides the foundation for developing values, principles and competencies for trauma-informed and developmentally sensitive health promotion.

# References

- Afifi, T.O., MacMillan, H.L., Boyle, M., Cheung, K., Taillieu, T., Turner, S., & Sareen, J. (2016). Child abuse and physical health in adulthood. *Health Reports, 27*(3), 10-18.
- Afifi, T.O., MacMillan, H.L., Boyle, M., Taillieu, T., Cheung, K., & Sareen, J. (2014). Child abuse and mental disorders in Canada. *CMAJ, Early Online*. DOI: 10.1503/cmaj.131792
- BC Provincial Mental Health and Substance Use Planning Council. (2013). *Trauma-Informed Practice Guide*. Vancouver, BC. Retrieved from [http://bccewh.bc.ca/wp-content/uploads/2012/05/2013\\_TIP-Guide.pdf](http://bccewh.bc.ca/wp-content/uploads/2012/05/2013_TIP-Guide.pdf)
- Beydoun, H. A., Beydoun, M. A., Kaufman, J. S., Lo, B., & Zonderman, A. B. (2012). Intimate partner violence against adult women and its association with major depressive disorder, depressive symptoms and postpartum depression: A systematic review and meta-analysis. *Social Science & Medicine, 75*(6), 959-975.
- Black, M.C., Basile, K.C., Breiding, M.J., Smith, S.G., Walters, M.L., Merrick, M.T., Chen, J. & Stevens, M. R. (2011). *The national intimate partner and sexual violence survey: 2010 summary report*. Atlanta: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
- Blasco-Ros, C., Herbert, J. & Martinez, M. (2014). Different profiles of mental and physical health and stress hormone response in women victims of intimate partner violence. *Journal of Acute Disease, 3*(4), 303-313.
- Briere, J., & Jordan, C. E. (2009). Childhood maltreatment, intervening variables, and adult psychological difficulties in women: An overview. *Trauma, Violence, & Abuse, 10*(4), 375-388.
- Bryant-Davis, T. (2010). Cultural considerations of trauma: Physical, mental, and social correlates of intimate partner violence exposure. *Psychological Trauma: Theory, Research, Practice, and Policy (Special Issue), 2*(4). 263-265.
- Canadian Population Health Initiative (2004). *Improving the health of Canadians*. Ottawa, ON: Canadian Institute for Health Information.
- Canadian Population Health Initiative. (2008). *Reducing gaps in health: A focus on socio-economic status in urban Canada*. Ottawa, ON: Canadian Institute for Health Information.
- Centers for Disease Control and Prevention (CDC). (2015). Intimate partner violence surveillance: Uniform definitions and recommended data elements. Retrieved from <http://www.cdc.gov/violenceprevention/pdf/intimatepartnerviolence.pdf>
- Centers for Disease Control and Prevention (CDC). (2015). The Social-Ecological Model: A Framework for Violence Prevention. Retrieved from [https://www.cdc.gov/violenceprevention/pdf/sem\\_framework-a.pdf](https://www.cdc.gov/violenceprevention/pdf/sem_framework-a.pdf)
- Clum, G. A., Nishith, P., & Resick, P. (2001). Traumarelated sleep disturbance and self-reported physical health symptoms in treatment-seeking female rape victims. *Journal of Nervous and Mental Disease, 189*(9), 618-622.
- Crenshaw, K. (1991). Mapping the margins: Intersectionality, identity politics, and violence against women of color. *Stanford Law Review, 43*(6), 1241-1299.
- D'Andrea, W., Sharma, R., Zelechowski, A., & Spinazzola, J. (2011). Physical health problems after single trauma exposure: When stress takes root in the body. *Journal of the American Psychiatric Nurses Association, 17*(6), 378-392.
- Dill, B. T., & Zambrana, R. E. (2009). Critical thinking about inequality: An emerging lens. In B. T. Dill & R. E. Zambrana (Eds.), *Emerging intersections: Race, class, and gender in theory, policy, and practice* (pp. 1-21). New Brunswick, NJ: Rutgers University Press.
- Dillon, G., Hussain, R., Loxton, D., & Rahman, S. (2013). Mental and physical health and intimate partner violence against women: A review of the literature. *International Journal of Family Medicine, 2013*, 1-16.

- Dutton, M. A., Green, B. L., Kaltman, S. I., Roesch, D.M., Zeffiro, T. A., & Krause, E. D. (2006). Review: Intimate partner violence, PTSD, and adverse health outcomes. *Journal of Interpersonal Violence, 21* (7), 955–968.
- Edleson, J. L. (1999). The overlap between child maltreatment and woman battering. *Violence Against Women, 5*(2), 134–154.
- Ely, G. E., Dulmus, C. N., & Wodarski, J. S. (2004). Domestic violence: A literature review reflecting an international crisis. *Stress, Trauma, and Crisis: An International Journal, 7*(2), 77-91.
- Elder, Jr., G. H., Johnson, M. K., & Crosnoe, R. (2003). The emergence and development of life course theory. In J.T. Mortimer and M.J. Shanahan (Eds.), *Handbook of the Life Course* (pp.3 -19). New York: Springer US, Kluwer Academic/Plenum Publishers.
- Engnes, K., Lidén, E., & Lundgren, I. (2012). Experiences of being exposed to intimate partner violence during pregnancy. *International Journal of Qualitative Studies on Health and Well-Being, 7*, 10.3402/qhw.v7i0.11199.
- Etherington, N. & Baker, L.L. (2016a). *The link between boys' victimization and adult perpetration of intimate partner violence: Opportunities for prevention across the life course*. London, Ontario: Centre for Research & Education on Violence Against Women & Children. ISBN 978-0-9688655-4-5
- Etherington, N. & Baker, L. (2016b). From “buzzword” to best practice: Applying intersectionality to children exposed to intimate partner violence. *Trauma, Violence & Abuse*, early online. DOI: 10.1177/1524838016631128
- Fry, D., McCoy, A., & Swales, D. (2012). The consequences of maltreatment on children’s lives: A systematic review of data from the East Asia and Pacific region. *Trauma, Violence, & Abuse, 13*(4), 209-233.
- Garcia-Moreno, C., Jansen, H A.F.M., Ellsberg, M., Heise, L., Watts, C. World Health Organization (WHO). (2005). *Multi-country study on women’s health and domestic violence against women*. Retrieved from <http://www.who.int/reproductivehealth/publications/violence/24159358X/en/>
- Government of Canada. (2008). *The Chief Public Health Officer’s report on the state of public health in Canada 2008*. Ottawa, ON: Public Health Agency of Canada.
- Hamby, S., Finkelhor, D., Turner, H., & Ormrod, R. (2010). The overlap of witnessing partner violence with child maltreatment and other victimizations in a nationally representative survey of youth. *Child Abuse and Neglect, 34*(10), 734–741.
- Hambrick, E. P., Tunno, A. M., Gabrielli, J., Jackson, Y., & Belz, C. (2014). Using multiple informants to assess child maltreatment: Concordance between case file and youth self-report. *Journal of Aggression, Maltreatment & Trauma, 23*(7), 751-771. Thousand Oaks, CA: Sage.
- Hankivsky, O. (2011). (Ed.). *Health inequities in Canada: Intersectional frameworks and practices*. Vancouver: UBC Press.
- Haskell, L. (2001). *Bridging responses: A front-line worker’s guide to supporting women who have post-traumatic stress*. Toronto, Ontario: Centre for Addiction and Mental Health.
- Health Canada. (2005). Child maltreatment: A public health issue. *Health Policy Research Bulletin, 9*, 1-40.
- Health Disparities Task Group of the Federal/Provincial/Territorial Advisory Committee on Population Health and Health Security. (2005). *Reducing health disparities – Roles of the health sector: Discussion Paper*. Ottawa, ON: Public Health Agency of Canada.
- Hebert, M., Cenat, J.M., Blais, M., Lavoie, F., Guerrier, M. (2016). Child Sexual Abuse, Bullying, Cyberbullying, and Mental health Problems Among High Schools Students: A Moderated Mediated Model. *Depression and Anxiety, 33* (7), 623-629.

- Herrenkohl, T.I., Hong, S., Klika, J.B., Herrenkohl, R.C., & Russo, M.J. (2013). Developmental impacts of child abuse and neglect related to adult mental health, substance use, and physical health. *Journal of Family Violence, 28*(2), 191-199.
- Herrenkohl, T. I., Sousa, C., Tajima, E. A., Herrenkohl, R. C., & Moylan, C. a. (2008). Intersection of child abuse and children's exposure to domestic violence. *Trauma, Violence & Abuse, 9* (2), 84-99.
- Herman, J. L. (1992a). Complex PTSD: A syndrome in survivors of prolonged and repeated trauma. *Journal of Traumatic Stress, 5* (3), 377-391.
- Herman, J. L. (1992b). *Trauma and Recovery: the Aftermath of Violence*. New York: Basic Books.
- Irish, L., Kobayashi, I., & Delahanty, D. L. (2010). Long-term physical health consequences of childhood sexual abuse: A meta-analytic review. *Journal of Pediatric Psychology, 35* (5), 450-461. jsp118.
- Jones, L., Hughes, M., & Unterstaller, U. (2001). Post-traumatic stress disorder (PTSD) in victims of domestic violence: A review of the research. *Trauma, Violence, & Abuse, 2*(2), 99-119.
- Jouriles, E. N., McDonald, R., Slep, A. M. S., Heyman, R. E., & Garrido, E. (2008). Child abuse in the context of domestic violence: prevalence, explanations, and practice implications. *Violence and Victims, 23*(2), 221-235.
- Katerndahl, D. A., Burge, S. K., Ferrer, R. L., Becho, J., & Wood, R. (2010). Complex dynamics in intimate partner violence: A time series study of 16 women. *Primary Care Companion to The Journal of Clinical Psychiatry, 12*(4), PCC.09m00859.
- Klinic Community Health Centre. (2013). *Trauma-informed: The trauma toolkit*. Retrieved from [http://trauma-informed.ca/wp-content/uploads/2013/10/Trauma-informed\\_Toolkit.pdf](http://trauma-informed.ca/wp-content/uploads/2013/10/Trauma-informed_Toolkit.pdf)
- Lagdon, S., Armour, C., & Stringer, M. (2014). Adult experience of mental health outcomes as a result of intimate partner violence victimisation: A systematic review. *European Journal of Psychotraumatology, 5* (24794), 1-12.
- Lind, Mackenzie J., Steven H. Aggen, Kenneth S. Kendler, Timothy P. York, and Ananda B. Amstadter. (2016). An epidemiologic study of childhood sexual abuse and adult sleep disturbances. *Psychological Trauma: Theory, Research, Practice, and Policy, 8* (2): 198-205.
- Lopez-Martinez, A.E., Serrano-Ibanez, E.R., Ruiz-Parraga, G.T., Gomez-Perez, L., Ramirez-Maestre C., & Esteve, R. (2016). Physical health consequences of interpersonal trauma: A systematic review of the role of psychological variables. *Trauma, Violence & Abuse*, doi: 10.1177/1524838016659488
- Lum On, M., Ayre, J., Webster, K., Moon, L., (2016). Examination of the health outcomes of intimate partner violence against women: State of knowledge paper. Sydney: Australia's National Research Organization for Women's Safety (ANROWS).
- Macy, R. J., Ferron, J., & Crosby, C. (2009). Partner violence and survivors' chronic health problems: Informing social work practice. *Social Work, 54*(1), 29-43.
- Magruder, K. M., Kassam-Adams, N., Thoresen, S., & Olf, M. (2016). Prevention and public health approaches to trauma and traumatic stress: a rationale and a call to action. *European Journal of Psychotraumatology, 1*, 1-9.
- Mason, S.M., Wright, R.J., Hibert, E.N., Spiegelman, D., Jun, H., Hu, F.B., & Rich-Edwards, J.W. (2013). Intimate partner violence and incidence of type 2 diabetes in women. *Diabetes Care, 36*(5), 1159-1165.
- Miller, E.C., Goodman, L.A., Thomas, K.A., Peterson, A., Scheer, J.R., Woulfe, J.M., & Warshaw, C. (2016). *Trauma-informed approaches for LGBTQ\* survivors of intimate partner violence: A review of literature and a set of practice observations*. Cambridge, MA: GLBTQ Domestic Violence Project.

- Miszkurka, M., Steensma, C., & Phillips, S.P. (2016). Correlates of partner and family violence among older Canadians: a life-course approach. *Health Promotion and Chronic Disease Prevention in Canada: Research, Policy and Practice*, 36(3), 45-53.
- Out, D., Hall, R.J., Granger, D.A., Page, G.G., & Woods, S. (2012). Assessing salivary C-reactive protein: Longitudinal associations with systemic inflammation and cardiovascular disease risk in women exposed to intimate partner violence. *Brain, Behavior, and Immunity*, 26(4), 543-551.
- Pan-Canadian Public Health Network. (2010). *Indicators of health inequalities: A report from the Population Health Promotion Expert Group and the Healthy Living Issue Group for the Pan-Canadian Public Health Network*. Ottawa, ON: Author.
- Pearlin, L. I. (2010). The life course and the stress process: Some conceptual comparisons. *Journals of Gerontology - Series B Psychological Sciences and Social Sciences*, 65 B(2), 207-215.
- Petrov, M., Davis, M., Belyea, M., & Zautra, A. (2016). Linking childhood abuse and hypertension: sleep disturbance and inflammation as mediators. *Journal of Behavioral Medicine*, 39(4), 716-726.
- Public Health Agency of Canada. (2016). Social determinants of health. Retrieved from <http://cbpp-pcpe.phac-aspc.gc.ca/public-health-topics/social-determinants-of-health/>
- Public Health Agency of Canada. (2014). What is family violence? Retrieved from <http://www.phac-aspc.gc.ca/sfv-avf/info/fv-eng.php>
- Public Health Agency of Canada. (2010). *Canadian incidence study of reported child abuse and neglect – 2008: Major findings*. Ottawa, ON: Author.
- Raphael, D. (2009). *Social Determinants of Health: Canadian Perspectives*, 2<sup>nd</sup> edition. Toronto: Canadian Scholars' Press.
- Reiche, E.M., Nunes, S.O., & Morimoto, H.K. (2004). Stress, depression, the immune system, and cancer. *Lancet Oncology*, 5 (10), 617-625.
- Rodgers, C. S., Lang, A. J., Laffaye, C., Satz, L. E., Dresselhaus, T. R., & Stein, M. B. (2004). The impact of individual forms of childhood maltreatment on health behavior. *Child Abuse & Neglect*, 28(5), 575-586.
- Rogosch, F. A., Dackis, M. N., & Cicchetti, D. (2011). Child maltreatment and allostatic load: consequences for physical and mental health in children from low-income families. *Development and Psychopathology*, 23(4), 1107-1124.
- Romano, E., Babchishin, L., Marquis, R., & Fréchette, S. (2015). Childhood maltreatment and educational outcomes. *Trauma, Violence, & Abuse*, 16(4), 418-437.
- SAMHSA. (2014a). *A treatment improvement protocol: Trauma-informed care in behavioral health services*. Rockville, MD: Department of Health and Human Services.
- SAMHSA (2014b). SAMHSA's Concept of Trauma and Guidance for a Trauma-Informed Approach. U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Office of Policy, Planning and Innovation. SMA 12-4884.
- Scott, S. & Babcock, J.C. (2010). Attachment as a moderator between intimate partner violence and PTSD symptoms. *Journal of Family Violence*, 25(1), 1-9.
- Seng, J. S., Graham-Bermann, S. A., Clark, K., McCarthy, A. M., & Ronis, D. L. (2005). Posttraumatic stress disorder and physical comorbidity among female children and adolescents: Results from service-use data. *Pediatrics*, 116 (6), 767-776.
- Shaw, B.A., & Krause, N. (2002). Exposure to physical violence during childhood, aging, and health. *Journal of Aging and Health*, 14(4), 467-494.

- Sinha, M. (2012). Section 2: Violence Against Intimate Partners. Statistics Canada.  
<http://www.statcan.gc.ca/pub/85-002-x/2012001/article/11643/11643-2-eng.htm>
- Sinha, M. (2013). Section 3: Intimate Partner Violence. Statistics Canada.  
<http://www.statcan.gc.ca/pub/85-002-x/2013001/article/11805/11805-3-eng.htm>
- Sinha, V., Trocme, N., Fallon, B., MacLaurin, B., Fast, E., Prokop, S. T., et al, (2011). Kiskisik Awasisak: Remember the Children. Understanding the Overrepresentation of First Nations Children in the Child Welfare System. Ontario: Assembly of First Nations.
- Sokoloff, N. J., & Pratt, C. (2005). *Domestic violence at the margins: Readings on race, class, gender, and culture*. New Jersey: Rutgers University Press.
- Tonmyr, L., Ouimet, C., & Ugnat, A. (2012). A Review of Findings from the Canadian Incidence Study of Reported Child Abuse and Neglect (CIS). *Canadian Journal of Public Health, 103*(2), 103-112.
- Van Ameringen, M., Mancini, C., Patterson, B., & Boyle, M. H. (2008). Post-Traumatic Stress Disorder in Canada. *CNS Neuroscience & Therapeutics, 14*(3), 171–181.
- Wathen, N. (2012). Health impacts of violent victimization on women and their children. *Department of Justice Canada*. Retrieved from [http://www.justice.gc.ca/eng/rp-pr/cj-jp/fv-vf/rr12\\_12/rr12\\_12.pdf](http://www.justice.gc.ca/eng/rp-pr/cj-jp/fv-vf/rr12_12/rr12_12.pdf)
- Williams, L. M. (2003). Understanding child abuse and violence against women: A life course perspective. *Journal of Interpersonal Violence, 18*(4), 441-451.
- Wong, J., & Mellor, D. (2014). Intimate partner violence and women's health and wellbeing: impacts, risk factors and responses. *Contemporary Nurse, 46*(2), 170-179.
- Woods, A. B., Page, G. G., O'Campo, P., Pugh, L. C., Ford, D., & Campbell, J.C. (2005). The mediation effect of posttraumatic stress disorder symptoms on the relationship between intimate partner violence and IFN-?? levels. *American Journal of Community Psychology, 36* (1-2), 159–175.
- World Health Organization (WHO), Department of Reproductive Health and Research, London School of Hygiene and Tropical Medicine, South African Medical Research Council. (2013). *Global and regional estimates of violence against women: Prevalence and health effects of intimate partner violence and non-partner sexual violence*. Retrieved from <http://www.who.int/reproductivehealth/publications/violence/9789241564625/en/>
- World Health Organization (WHO). (2016). Violence and injury prevention: Child maltreatment. Retrieved from [http://www.who.int/violence\\_injury\\_prevention/violence/child/en/](http://www.who.int/violence_injury_prevention/violence/child/en/)
- Zimmerman, F., Mercy, J. A. (2010). *A Better Start: Child Maltreatment Prevention as a Public Health Priority*. Washington, D.C.: ZERO TO THREE: National Center for Infants, Toddlers, and Families. Retrieved: [https://vetoviolence.cdc.gov/apps/phl/docs/A\\_Better\\_Start.pdf](https://vetoviolence.cdc.gov/apps/phl/docs/A_Better_Start.pdf)